

# Underage Drinking in Montana


## *The Facts*

Tragic health, social, and economic problems result from the use of alcohol by youth. Underage drinking is a causal factor in a host of serious problems, including homicide, suicide, traumatic injury, drowning, burns, violent and property crime, high risk sex, fetal alcohol syndrome, alcohol poisoning, and need for treatment for alcohol abuse and dependence.

### *Problems and Costs Associated with Underage Drinking in Montana*

Underage drinking cost the citizens of Montana \$0.2 billion in 2010. These costs include medical care, work loss, and pain and suffering associated with the multiple problems resulting from the use of alcohol by youth.<sup>1</sup> This translates to a cost of \$2,580 per year for each youth in the State or \$3.04 per drink consumed underage. Excluding pain and suffering from these costs, the direct costs of underage drinking incurred through medical care and loss of work cost Montana \$107 million each year or \$1.33 per drink. In contrast, a drink in Montana retails for \$1.13.

**Costs of Underage Drinking  
Montana, 2010 \$**



**Total: \$0.2 billion**

**Costs of Underage Drinking by Problem, Montana, 2010 \$**

Problem	Total Costs (in millions)
Youth Violence	\$94.5
Youth Traffic Crashes	\$81.4
High-Risk Sex, Ages 14-20	\$18.1
Youth Property Crime	\$12.9
Youth Injury	\$10.6
Poisonings and Psychoses	\$2.4
FAS Among Mothers Age 15-20	\$5.7
Youth Alcohol Treatment	\$19.6
<b>Total</b>	<b>\$245.2</b>

Youth violence (homicide, suicide, aggravated assault) and traffic crashes attributable to alcohol use by underage youth in Montana represent the largest costs for the State. However, a host of other problems contribute substantially to the overall cost. Among teen mothers, fetal alcohol syndrome (FAS) alone costs Montana \$6 million.

Young people who begin drinking before age 15 are four times more likely to develop alcohol dependence

and are two and a half times more likely to become abusers of alcohol than those who begin drinking at age 21.<sup>2</sup> In 2009, 488 youth 12- 20 years old were admitted for alcohol treatment in Montana, accounting for 11% of all treatment admissions for alcohol abuse in the state.<sup>3</sup>

## ***Alcohol Consumption by Youth in Montana***

Underage drinking is widespread in Montana. Approximately 44,000 underage customers in Montana drink each year. In 2009, Montana students in grades 9-12 reported:<sup>4</sup>

- 75.7% had at least one drink of alcohol on one or more days during their life.
- 24.0% had their first drink of alcohol, other than a few sips, before age 13.
- 42.8% had at least one drink of alcohol on one or more occasion in the past 30 days.
- 30.1% had five or more drinks of alcohol in a row (binge drinking) in the past 30 days.
- 5.1% had at least one drink of alcohol on school property in the past 30 days.

In 2009, underage customers consumed 23.9% of all alcohol sold in Montana, totaling \$91 million in sales (in 2010 dollars). These sales provided profits of \$45 million to the alcohol industry.<sup>1</sup> Ranking states based on the percentage of alcohol consumed underage, with 1 the highest, Montana ranked number 23. This percentage is affected by both adult and youth drinking levels.

Annual sales of alcohol consumed by youth in Montana averaged \$2,075 per underage customer. Underage customers were heavier consumers than adults. They drank an average of 5.0 drinks per day; in contrast, legal customers consumed only 2.0.

## ***Harm Associated with Underage Drinking in Montana***

Underage drinking in Montana leads to substantial harm due to traffic crashes, violent crime, property crime, unintentional injury, and risky sex.

- During 2009, an estimated 10 traffic fatalities and 470 nonfatal traffic injuries were attributable to driving after underage drinking.
- In 2009, an estimated 3 homicides; 2,300 nonfatal violent crimes such as rape, robbery and assault; and 4,900 property crimes including burglary, larceny, and car theft were attributable to underage drinking.
- In 2007, an estimated 2 alcohol involved fatal burns, drownings, and suicides were attributable to underage drinking.
- In 2009, an estimated 130 teen pregnancies and 2,799 teens having risky sex were attributable to underage drinking.

For comparison with other states, in US rather than state prices, the harm from underage drinking per youth in Montana averages \$1,122. Such comparisons require caution. In part, they may reflect differences in crime and crash rates, problem-reporting to police, and co-occurring drug use.

Produced by the Pacific Institute for Research and Evaluation (PIRE) with funding from the Office of Juvenile Justice and Delinquency Prevention (OJJDP), September 2011.

---

<sup>1</sup> Levy, D.T., Miller, T.R., & Cox, K.C. (2003). Underage drinking: societal costs and seller profits. Working Paper. Calverton, MD: PIRE.

<sup>2</sup> Grant, B.F., & Dawson, D.A. (1997). Age at onset of alcohol use and its association with DSM-IV alcohol abuse and dependence: Results from the National Longitudinal Alcohol Epidemiologic Survey. *Journal of Substance Abuse* 9: 103-110.

<sup>3</sup> Office of Applied Studies, Substance Abuse and Mental Health Services Administration. Treatment Episode Data Set. (2011). *Substance Abuse Treatment by Primary Substance of Abuse, According to Sex, Age, Race, and Ethnicity, 2009*. Available [On-line]: <http://www.icpsr.umich.edu/icpsrweb/SAMHDA/studies/30462>.

<sup>4</sup> Center for Disease Control (CDC). (2011). Youth Risk Behavior Surveillance System (YRBSS). Available [On-line]: <http://apps.nccd.cdc.gov/youthonline/App/Default.aspx>.